

rane mixtures: a solution- state  $^{11}\text{B}$  NMR study. *Phys. Chem. Chem. Phys.*, 2011;13:3809–18.

[38] Janda R., Heller G. IR- and Ramanspektren isotope markierter tetra- und pentaborate. *Spectrochim Acta*, 1982;36A:997–1001.

[39] Jun L., Shuping X., Shiyang G. FT-IR and Raman spectroscopic study of hydrated borates. *Spectrochim Acta*, 1995;51:519–32.

[40] Demirci U.B., Bernard S., Chiriach R., Toche F., Miele P. Hydrogen release by thermolysis of ammonia borane  $\text{NH}_3\text{BH}_3$  and then hydrolysis of its by-product  $[\text{BNH}_x]$ . *J. Power Sources*, 2011;196:279–86.

[41] Singh S.K., Xu Q. Bimetallic Ni-Pt nanocatalysts for selective decomposition of hydrazine in aqueous solution to hydrogen at room temperature for chemical hydrogen storage. *Inorg. Chem.*, 2010;49:6148–52.

[42] Zhu Q.L., Zhong D.C., Demirci U.B., Xu Q. Controlled synthesis of ultrafine surfactant-free NiPt nanocatalysts toward efficient and complete hydrogen generation from hydrazine borane at room temperature. *ACS Catal.*, 2014;4:4261–9.

[43] Zahmakıran M., Özkır S. Transition metal nanoparticles in catalysis for the hydrogen generation from the hydrolysis of ammonia-borane. *Top Catal.*, 2013;56:1171–83.

[44] Patel N., Miotello A. Progress in Co-B related catalyst for hydrogen production by hydrolysis of boronhydrides: a review and the perspectives to substitute noble metals. *Int. J. Hydrogen Energy*, 2015;40:1429–64.


[45] Gao W., Li C., Wu M., He S., Wei M., Evans D.G., et al. Supported nickel-iron nanocomposites as a bifunctional catalyst towards hydrogen generation from  $\text{N}_2\text{H}_4\cdot\text{H}_2\text{O}$ . *Green Chem.*, 2014;16:1560–8.

[46] Singh S.K., Xu Q. Complete conversion of hydrous hydrazine to hydrogen at room temperature for chemical hydrogen storage. *J. Am. Chem. Soc.*, 2009;131:18032–3.

[47] Singh S.K., Xu Q. Bimetallic nickel-iridium nanocatalysts for hydrogen generation by decomposition of hydrous hydrazine. *Chem. Commun.*, 2010;46:6545–7.

[48] Singh S.K., Iizuka Y., Xu Q. Nickel-palladium nanoparticle catalyzed hydrogen generation from hydrous hydrazine for chemical hydrogen storage. *Int. J. Hydrogen Energy*, 2011;36:11794–801.

[49] Schlesinger H.I., Burg A.B. Hydrides of boron. I. An efficient new method of preparing diborane; new reactions for preparing bromo-diborane and the stabler penta-borane,  $\text{B}_5\text{H}_9$ . *J. Am. Chem. Soc.*, 1931;53:4321–32.



Dear Colleagues,

On behalf of the Organizing Committee, we are pleased to invite you to take part in the “International Conference on Nanomaterials, Nanofabrication and Nanocharacterization (NANOMACH)” which will be held on April 23-29, 2020 in the stunning Convention Centre of one of the largest international resort in Turkey right in the heart of Blue Lagoon (Oludeniz) area.

NANOMACH is traditional annual congress for scientists and expected to collect a wide audience of participants and listeners. We are planning an outstanding scientific program led by world-renowned invited speakers, that will not only be a platform to showcase exciting new developments in nanotechnology, but will also reveal the transformational role of nanotechnology in supporting a range of physical and life sciences. We hope that the scientific program, including a wide array of topics will live to your expectations, and that participation in the congress will offer you an opportunity to meet up with your colleagues, friends and renowned specialists from all over the world.

We look forward to seeing you in Fethiye.

Organizing Committee

<http://www.nanomach.org/Pages/topics>

