

## DIRECT RADIO PROBING AND INTERPRETATION OF THE SUN'S DENSITY PROFILE

IVER H. CAIRNS<sup>1</sup>, Vasili V. Lobzin<sup>1</sup>, Alexander Warmuth<sup>2</sup>, Bo Li<sup>1</sup>, Peter A. Robinson<sup>1</sup>, and Gottfried Mann<sup>2</sup>

<sup>1</sup>School of Physics, University of Sydney, NSW 2006, Australia

<sup>2</sup>Astrophysikalisches Institut Potsdam, An der Sternwarte 16, D 14482 Potsdam, Germany

The Sun's electron number density profile  $n_e(r)$  is not well measured and understood from the Sun's photosphere ( $r = R_S$ ) to the solar wind (say,  $r > 10 R_S$ ), despite its great importance for solar physics. A powerful new technique is developed and applied to extract  $n_e(r)$  directly from the time-varying frequency of coronal type III radio bursts, for the first time. Unexpectedly, wind-like regions with  $n_e = A (r - R_S)^{-2}$  are common below  $2R_S$ , and coronal type III bursts often are almost linear in  $1/f - t$  dynamic spectra. Indices lower and higher than 2 also occur, the former plausibly due to non-radial curved magnetic field lines. The profile  $n_e = A (r - R_S)^{-2}$  is consistent with the coronal radio data, type III simulations, and previous models based on eclipse and coronagraph observations. It gives a first physical basis for the previous models. A simple interpretation is in terms of conical flow from localized sources close to the photosphere, plausibly corresponding to UV funnels. The results also imply that solar wind acceleration occurs where  $2 < r/R_S < 10$ .

solar radio bursts, solar corona, electron density profile

Iver H. Cairns, School of Physics, University of Sydney, NSW 2006, Australia. Ph +61-2-9351-3961, Fax +61-2-9351-7726, Email [i.cairns@physics.usyd.edu.au](mailto:i.cairns@physics.usyd.edu.au) .