

DYNAMIC EVOLUTION OF THE RADIATION BELT PHASE SPACE DENSITY

YURI SHPRITS¹, Bibin Ni¹, Reiner Friede², Yue Chen², Dmitri Kondrashov¹, Dmitri Subbotin¹, Geoff Reeves², Michael Ghil¹, Richard Thorne¹, Tsugunobu Nagai³

¹Department of Atmospheric and Oceanic Sciences,
UCLA, Los Angeles, CA 90095, USA

²Los Alamos National Laboratory
Los Alamos, New Mexico, 87545, USA

³Tokyo Institute of Technology
Earth and Planetary Sciences I 2-5
Tokyo 152-8551, Japan

We present radiation belt relativistic electron Phase Space Density (PSD) is obtained using the data assimilative VERB code combined with observations from GEO, CRRES, and Akebono data. Reanalysis of data shows the pronounced peaks in the phase space density and pronounced drop outs of fluxes during the main phase of a storm. Reanalysis results clearly show that persistent peaks in PSD independent of the assumed magnetic field model. Simulations with the 3D VERB code accounting for the violations of all three adiabatic invariants is presented. The results of the reanalysis are discussed and compared to the simulations with the recently developed VERB 3D code.

Keywords wave-particle interactions

Yuri Y Shprits, DAOS UCLA, Los Angeles, CA 90095, USA.

yshprits@atmos.ucla.edu