

PHASE RELATION BETWEEN PI 2-ASSOCIATED IONOSPHERIC DOPPLER VELOCITY AND MAGNETIC PULSATION AT MID-LATITUDE MAGDAS STATION

AKIHIRO IKEDA¹, Kiyohumi Yumoto², Teiji Uozumi², Manabu Shinohara³, Kenro Nozaki⁴, Akimasa Yoshikawa¹, V.V. Bychkov⁵, B. M. Shevtsov⁵

¹Department of Earth and Planetary Sciences, Kyushu University, Fukuoka, Japan

²Space Environment Research Center, Kyushu University, Fukuoka, Japan

³Kagoshima National College of Technology, Kagoshima, Japan

⁴National Institute of Information and Communications Technology, Tokyo, Japan

⁵IKIR of the Far Eastern Branch of Russian Academy of Sciences, Russia

Pi 2 magnetic pulsation, which is an impulsive hydromagnetic oscillation, occurs globally in the magnetosphere at the onset of magnetospheric substorms [e.g., *Saito and Matsushita*, 1968]. Pi 2 pulsations have various modes depending on latitude and local time [e.g., *Yumoto and the CPMN Group*, 2001]. In particular low- and mid-latitude Pi 2 pulsations are explained by the theory of the cavity mode resonance in the plasmasphere [e.g., *Sutcliffe and Yumoto*, 1991]

However, the examination of the Pi 2-associated Doppler velocity is limited. In the present paper, we examine the phase relation between the ionospheric Doppler velocity (V^*) in the ionospheric F-region detected by an FM-CW (Frequency Modulated Continuous Wave) radar and the ground magnetic H component (H) observed by MAGDAS magnetometer at PTK (L=2.05). In addition, we estimate the ionospheric electric field intensity.

From statistics, 114 Pi 2 events were observed during a 43-day period from 23 September 2006 to 4 November 2006. The variations of H and V^* revealed high coherence (correlation coefficient: $\gamma \geq 0.6$) for a half of the 114 Pi 2 events, for about a half of which the H and V^* variations had the same dominant frequency. For such events, the phase difference between H and V^* is -90° or 270° in the midnight sector (21-03 LT), and the averaged E_y (east-west electric field) amplitude derived from V^* is 0.24 mV/m. In contrast, no clear phase relation of H and V^* could be found for Pi 2 oscillations in the other local time sectors. Based on a box model of the ideal cavity mode by Takahashi et al. [2001], the phase relation of H and V^* for Pi 2s in the midnight sector (21-03 LT) may be explained in terms of the radial standing structure of compressional waves, i.e., cavity mode oscillation.

Pi2, Doppler Radar, Ionospheric Electric Field

Akihiro Ikeda, Department of Earth and Planetary Sciences, Kyushu University, Fukuoka, Japan
e-mail: a-ikeda@geo.kyushu-u.ac.jp