

THEMIS MULTI-SPACECRAFT OBSERVATIONS OF MAGNETOSHEATH PLASMA PENETRATION DEEP INTO THE LOW-LATITUDE DAYSIDE MAGNETOSPHERE DURING NORTHWARD IMF

MARIT ØIEROSET 1, Tai Phan 1, Vassilis Angelopoulos 2, Wenhui Li 3, Jimmy Raeder 3, Jonathan Eastwood 1, Jim McFadden 1, Davin Larson 1, Chuck Carlson 1, Karl-Heinz Glassmeier 4

1. Space Sciences Laboratory, University of California, Berkeley, CA 94720, USA, e-mail: oieroset@ssl.berkeley.edu

2. Institute for Geophysical and Planetary Physics, University of California, Los Angeles, California, USA

3. Space Science Center, University of New Hampshire, Durham, New Hampshire, USA

4. Institut für Geophysik und Extraterrestrische Physik, Technische Universität Braunschweig, Germany

During the THEMIS coast phase in 2007 and the dayside phase in 2008 magnetosheath plasma was often observed deep inside the magnetosphere when the IMF was northward directed. Using THEMIS multi-spacecraft observations we are able to determine the thickness and evolution of this layer. We present dayside boundary layer observations at different local times and for different IMF clock angles and compare with global MHD simulations to shed light on the entry process. This dayside boundary layer could be a source of the so-called cold dense plasma sheet (CDPS) frequently observed in the tail during northward IMF.

magnetopause, boundary layers

Marit Oieroset, Space Sciences Laboratory, University of California, 7 Gauss Way, Berkeley, 94720, USA, tel: +1 510 642 2290, e-mail: oieroset@ssl.berkeley.edu