

NONLINEAR MECHANISMS OF LOWER BAND AND UPPER BAND VLF CHORUS EMISSIONS IN THE MAGNETOSPHERE

YOSHIHARU OMURA ¹, Mitsuru Hikishima ^{1 2}, Yuto Katoh ^{1 3}, Danny Summers ^{1 4 5}, Satoshi Yagitani ²

1. Research Institute for Sustainable Humanosphere, Kyoto University, Kyoto Japan, e-mail: omura@rish.kyoto-u.ac.jp
2. Graduate School of Natural Science and Technology, Kanazawa University, Kanazawa, Japan.
3. Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, Miyagi, Japan.
4. Department of Mathematics and Statistics, Memorial University of Newfoundland, St. John's, Newfoundland, Canada.
5. School of Space Research, Kyung Hee University, Yongin, Gyeonggi, Korea.

We develop a nonlinear wave growth theory of magnetospheric chorus emissions, taking into account the spatial inhomogeneity of the static magnetic field and the plasma density variation along the magnetic field line. We derive theoretical expressions for the nonlinear growth rate and the amplitude threshold for the generation of self-sustaining chorus emissions. We assume that nonlinear growth of a whistler-mode wave is initiated at the magnetic equator where the linear growth rate maximizes. Self-sustaining emissions become possible when the wave propagates away from the equator during which the increasing gradients of the static magnetic field and electron density provide the conditions for nonlinear growth. The amplitude threshold is tested against both observational data and self-consistent particle simulations of the chorus emissions. The self-sustaining mechanism can result in a rising tone emission covering the frequency range of 0.1 - 0.7 of the equatorial electron gyrofrequency. During propagation higher frequencies are subject to stronger dispersion effects that can destroy the self-sustaining mechanism. We obtain a pair of coupled differential equations for the wave amplitude and frequency. Solving the equations numerically, we reproduce a rising tone of VLF whistler-mode emissions that is continuous in frequency. Chorus emissions, however, characteristically occur in two distinct frequency ranges, a lower band and an upper band, separated at half the electron gyrofrequency. We explain the gap by means of the nonlinear damping of the longitudinal component of a slightly oblique whistler-mode wave packet propagating along the inhomogeneous static magnetic field.

whistler-mode wave, chorus, relativistic electrons

Yoshiharu Omura, Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan, tel: 81-774-8463, fax: 81-774-31-8463, e-mail omura@rish.kyoto-u.ac.jp