

# **THE EFFECT OF IONOSPHERIC PEDERSEN CONDUCTANCE ON THE FORMATION OF VORTICES IN THE MAGNETOTAIL**

RAYMOND J. WALKER 1, Maha Ashour Abdalla 2 and Mostafa El-Alaoui 3

1. Institute of Geophysics and Planetary Physics and Department of Earth and Space Sciences, University of California, Los Angeles, CA. 90095-1567 U.S.A. email: [rwalker@igpp.ucla.edu](mailto:rwalker@igpp.ucla.edu)

2. Institute of Geophysics and Planetary Physics and Department of Physics and Astronomy, University of California, Los Angeles, CA 90095-1567 U.S. A. email: [mabdalla@igpp.ucla.edu](mailto:mabdalla@igpp.ucla.edu)

3. Institute of Geophysics and Planetary Physics, University of California, Los Angeles, CA 90095-1567 U. S. A. email: [mostafa@igpp.ucla.edu](mailto:mostafa@igpp.ucla.edu).

In a series of simulation studies of the magnetosphere during prolonged intervals when the interplanetary magnetic field (IMF) was southward we found that earthward convection from a tail neutral line could stop and reverse direction forming large vortices in the tail. We have investigated the effects of ionospheric Pedersen conductance on magnetospheric convection by carrying out a series of simulations with steady southward IMF and fixed ionospheric conductance. For small conductances (3S and 6S) a “classical” convection pattern in which a near-Earth neutral line returns reconnected magnetic flux to the dayside occurs. However, for higher conductance (20S) like that found in the auroral zone during substorms flow in the near-Earth tail became stagnant and then vortices formed. We show that pressure gradients in the near-Earth plasma sheet alone are insufficient to cause the formation of the vortices and that a combination of ionospheric line tying and pressure gradients is needed.

Magnetospheric convection, vorticity, ionospheric conductance

Raymond J. Walker, IGPP. UCLA Los Angeles, CA 90095-1567, U.S. A.