

MOMENTUM TRANSFER ON OLD OPEN FIELD LINES AND THE ROLES OF THE IMF EAST-WEST COMPONENT (BY) AND CONDUCTIVITY GRADIENTS

Per Even Sandhøolt (1), Charles J. Farrugia (2)

(1) Department of Physics, University of Oslo, Sem Sælands vei 24, Blindern, Oslo, 0316, Norway.

(2) Space Science Center, University of New Hampshire, 39 College Road, Durham, NH 03824, United States

Using DMSP F13 data we investigate the newly discovered channels of enhanced antisunward convection occurring at the dawn (0600-0900 MLT) or dusk (1500-1800 MLT) flanks of the polar cap for the different combinations of IMF By polarity, hemisphere and the dawn/dusk sides. Dawn-side cases occur for the following combinations of hemisphere (NH/SH) and By polarity: NH-dawn/By > 0 and SH-dawn/By < 0. The dusk-side cases are: NH-dusk/By < 0 and SH-dusk/By > 0 .

The flow channels are placed in the context of particle precipitation regimes/boundaries and ionospheric conductivity gradients. They are found to be threaded by "old open field lines" characterized by polar rain precipitation.

In the dawn-side cases (NH-dawn/By > 0 and SH-dawn/By < 0) the polar rain contains the "solar wind strahl" component. The convection enhancement is attributed to the Pedersen current closure of Birkeland currents poleward of the R1/R2 currents.

This is momentum transfer from the solar wind via dynamo action at the high-latitude boundary layer (HBL). The conductivity gradient at the polar cap boundary contributes to establishing the convection channel and the enhancement of the dawn-dusk convection asymmetry extending beyond the dawn-dusk terminator during intervals of nonzero IMF By component. The HBL - ionosphere coupling via Birkeland currents is a source of dawn-dusk convection asymmetry and Svalgaard - Mansurov effect which comes in addition to the effect of magnetic tension acting on "newly open field lines".

Momentum, conductivity

Per Even Sandhøolt, Department of Physics, University of Oslo, Sem Sælands vei 24, Blindern, Oslo, 0316, Norway.