

FLOW, AURORA AND PI2 ASSOCIATIONS OBSERVED BY THEMIS

LARRY KEPKO 1, Emma Spanswick 2, Eric Donovan 2, Vassilis Angelopoulos 3, Joachim Raeder 1, Jim McFadden 4, K.-H. Glassmeier 5

Space Science Center, University of New Hampshire, Durham, NH, USA, e-mail:

larry.kepko@unh.edu, j.raeder@unh.edu

University of Calgary, Calgary, AB, Canada, e-mail: emma@phys.ucalgary.ca,

edonovan@ucalgary.ca

University of California, Los Angeles, Los Angeles, CA, USA, e-mail: vassilis@ucla.edu

University of California, Berkeley, Berkeley, CA, USA, e-mail:

mcfadden@ssl.berkeley.edu

Technical University of Braunschweig, Braunschweig, Germany, e-mail:

kh.glassmeier@tu-bs.de

It has been known for decades that auroral substorm onset occurs on (or near) the most equatorward growth-phase auroral arc, which is thought to map to the near-geosynchronous region. The lack of auroral signatures poleward of this arc prior to onset has been a major criticism of flow-burst driven models of substorm onset. The combined THEMIS 5 spacecraft in-situ and ground array measurements provide an unprecedented opportunity to examine the causal relationship between midtail plasma flows, aurora, and ground magnetic signatures. In this paper we present an event from 2008 using multi-spectral all sky imager data from Gillam and in-situ data from THEMIS. The multispectral data indicate an equatorward moving auroral form prior to substorm onset. When this forms reaches the most equatorward arc, the arc brightens and an auroral substorm begins. The THEMIS data show fast Earthward flows prior to onset as well. We suggest that the results strongly support flow-burst driven models of magnetospheric activity.

Substorm, THEMIS, Pi2, aurora

Larry Kepko, Space Science Center, University of New Hampshire, 8 College Road, Durham, NH, 03824, USA, tel: +603-862-3891, e-mail: larry.kepko@unh.edu