

LANL* V1.0: RADIATION BELT DRIFT SHELL MODELING FOR REAL-TIME AND REANALYSIS APPLICATIONS

JOSEF KOLLER 1, Geoffrey D. Reeves 1, Reiner H. W. Friedel 1

Los Alamos National Lab, USA, email: jkoller@lanl.gov

Radiation belt modeling requires detailed information about the particle drift shells and their spatial and temporal variations. The drift shells are commonly represented by the third adiabatic invariant, L^* . Calculating drift shells in realistic geomagnetic fields are very time consuming and typically involve three-dimensional numerical integration of the global field. Choosing a simplistic but computationally less expensive geomagnetic field model could lead to large inaccuracies and even wrong conclusions [Huang et al., 2008]. Efficient algorithms without the sacrifice of numerical accuracy are required especially for real-time radiation belt modeling and forecasting. Likewise, long-duration calculations (e.g. solar-cycle scales) are impractical with the standard methods. We have developed a new method for calculating accurate L^* values. Our technique is based on a neural network that is 5-6 orders of magnitude faster than the standard numerical integration and interpolation. Our technique holds promise to remove a major roadblock in radiation belt research. While the technique is applicable for any closed drift shell, we will present results of a detailed validation at geosynchronous orbit. The neural network is freely available as LANL* V1.0 library. The FORTRAN library is complete with examples, Makefile for IDL and FORTRAN, and available upon request.

Geomagnetic field, drift shell modeling, adiabatic invariants

Josef Koller, Los Alamos National Lab, NM 87544, USA, tel: +1 505 665 3399;
fax: +1 505 665 7395; email: jkoller@lanl.gov